RISK OF DEATH | VACCINATED VS UNVACCINATED

Increased Risk of Death Due to Vaccination Unvaccinated > 3x Hep B vaccine > 5x Pneumococcal vaccine > 6x Meningococcal vaccine > 9x Influenza vaccine > 12x DTP vaccine > 23x Hep A vaccine > 33x Hib vaccine >160x Varicella vaccine > 261x MMR vaccine > 4.600.000x Polio vaccine 50 100 150 200 250 300 4,000,000



IN EACH CASE, THE RISK HAS BEEN FOUND TO BE HIGHER FROM VACCINATION THAN FROM NON-VACCINATION



To Vaccinate or Not to Vaccinate?

Vaccines are promoted as "safe and effective", but no government agency has conducted a comprehensive analysis proving any vaccine to be of overall benefit. Benefit is assumed based on vaccine-induced antibody levels and minimal safety testing. Low incidence of any targeted disease and the avoidability of harm from it in the unvaccinated are also not taken into account.

To make an informed health care decision, the question every parent needs answered is –

WHICH CHOICE IS SAFER?

Recently, a mathematician using the government's own data compared the risk of death or serious condition from receipt, versus not-receipt, of each vaccination mandated for enrolment in U.S. daycare or educational institutions.

The results were that in each case, the risk was found to be higher from vaccination than from non-vaccination.



These Results Should Cause Every Parent To Pause For more details: https://vaccinechoicecanada.com/RiskOfDeath

