

compares BC Vaccine Schedules

www.vaccinechoicecanada.com

1 000			0015	
TOTAL PreSchool 22 Vaccines In 6 Injections & 4 oral doses		AGE Prenatal	TOTAL PreSchool 50 (53)* Vaccines In 19 (22)* Injections & 2 oral doses *Aboriginal babies receive 3 doses of Hep A	
Ø	Number Vaccines	4	Number Vaccines 1	Influenza Vaccine in routine schedule for pregnant women
DPT: 3 vaccines in 1 shot OPV: oral polio, live virus vaccine	4	2 Months	9	DTaP-HB-IPV-Hib 6 vaccines in 1 shot Meningcoccal C-C Pneumococcal C-13 Rotavirus (oral) live virus vaccine
DPT: 3 vaccines in 1 shot OPV: oral polio live virus vaccine	4	4 Months	8	DTaP-HB-IPV-Hib 6 vaccines in 1 shot Pneumococcal C-13 Rotavirus (oral) live virus vaccine
DPT: 3 vaccines in 1 shot	3	6 Months	8 (9)	DTaP-HB-IPV-Hib 6 vaccines in 1 shot Influenza 2 doses 4 weeks apart Hepatitis A*
Ø		12 Months	6	MMR 3 live virus vaccines in 1 shot Meningcoccal C-C
MMR: 3 live virus vaccines in 1 shot at 15 months	3	18 Months		Pneumococcal C-13 Varicella chickenpox live virus vaccine
DPT: 3 vaccines in 1 shot OPV: oral polio live virus vaccine	4	18 Months	6 (7)	DTaP-IPV-Hib 5 vaccines in 1 shot Influenza (1st Annual dose) Hep A*
Ø		3 Yr	1	Influenza (Annual dose in Fall)
DPT: 3 vaccines in 1 shot OPV: oral polio live virus vaccine	4 Total 22	4-6 Yr	11 (12) Total 50 (53)	Tdap-IPV 4 vaccines in 1 shot MMRV 4 live virus vaccines in 1 shot Influenza (3 Annual doses, 1 each Fall) Hep A*

By the time your child graduates from high school, they will have received an additional 11 annual Influenza vaccines, 2 doses *HPV in Grade 6, Tdap (3-in-1 shot) & Men C-ACYW-135 in Grade 9. *Ages 9–18 inclusive: All Girls (HPV-9). Boys (HPV-4) who are gay or questioning their sexual identity, HIV positive, street involved, in the custody of Ministry of Children & Families or in youth detention centres also recieve 2 doses of HPV.

TOTAL Grade School & High School in 2017 Girls: 17 Vaccines in 15 Injections most Boys: 15 Vaccines in 13 Injections

2017 GRAND TOTAL: Prenatal to Age 18 65–70 Vaccines

Did you know vaccination is VOLUNTARY in Canada?

As Health Canada explained years ago: "Unlike some countries, immunization is not mandatory in Canada; it cannot be made mandatory because of the Canadian Constitution...legislation and regulations must not be interpreted to imply compulsory immunization."

LEGEND

Vaccines used in the 1983 BC schedule

Note: All 3 of these vaccines have since been withdrawn from the market due to safety concerns.

DPT: diphtheria, whole cell pertussis, tetanus vaccine 3 vaccine combination

OPV: oral, live virus polio vaccine

MMR: measles, mumps (Urabe strain), rubella vaccine

3 live virus vaccine combination

Vaccines used in the 2017 BC schedule

DTaP-HB-IPV-Hib 6 vaccine combination with diphtheria(D), tetanus(T), acellular pertussis(aP), HepatitusB (HB), inactivated polio (IPV) & Hib (Haemophilus Influenzae B)

• INFANRIX hexa®

DTaP-IPV-Hib 5 vaccine combination with diphtheria(D), tetanus(T), acellular pertussis(aP), inactivated polio (IPV) & Hib (Haemophilus Influenzae B) vaccines

• PediacelTM or Infanrix-IPV/Hib[®]

Hepatitis A (HepA or HA) virus vaccine (Aboriginal only)

• AVAXIM® Pediatric, HAVRIX® 720 JUNIOR, VAQTA®

Hepatitis B (HB): genetically engineered virus vaccine

• ENGERIX®-B or RECOMBIVAX HB® pediatric

HPV Human papillomavirus vaccine,

• GARDASIL 9TM (girls), GARDASILTM (boys)

Influenza Annual Flu shot, quadrivalent vaccine recommended

• FLULAVAL®Tetra for babies 6 -23 months

 \bullet FLUMIST® Quadrivalent or FLULAVAL® Tetra 2 –17 yrs (If a quadrivalent product is unavailable, Fluviral® or Agriflu® used.)

Men C-C: Meningococcal serotype C vaccine,

• MENJUGATE® or NEISVAC-C®

Men C-ACYW-135: 4 Meningcococcal serotypes,

• MENVEO® or MENACTRA®

*MMR: measles, mumps, rubella,

3 combination live virus vaccine

• $MMRII^{\mathbb{R}}$ or $Priorex^{\mathbb{R}}$ for children < 4 years old

*MMRV: measles, mumps, rubella, & varicella (chickenpox) 4 combination live virus vaccine

• Priorex Tetra[®] or ProQuadTM for children ≥ 4 yrs old

Pneu C-13: pneumococcal vaccine, 13 serotypes,

• PREVNAR 13®

*Rotavirus: live virus oral vaccine,

• RotarixTM (2 doses) or RotaTeq® (3 doses)

Tdap-IPV: tetanus, diphtheria, acellular pertussis, polio 4 vaccine combination, age 4-6

• ADACEL-Polio® or BOOSTRIX-Polio®

Tdap: tetanus, diphtheria, acellular pertussis, polio 4 vaccine combination, Grade 9 booster

• ADACEL® or BOOSTRIX®

*Varicella: chickenpox live virus vaccine

•VARILRIX® or VARIVAX® III

*Live virus vaccines can shed for varying amounts of time (weeks to months) in the body fluids of vaccinated people and can be transmitted to others. "Little is known about the potential of live attenuated and genetically engineered vaccine viruses to mutate and recombine with other viruses and create new viruses that will cause disease or affect the integrity of the human genome, human microbiome and healthy functioning of the immune and neurological systems." NVIC http://www.nvic.org/cmstemplates/nvic/pdf/live-virus-vaccines-and-vaccine-shedding.pdf

What's In those Vaccines?

In the first year of life, during critical phases of brain and immune system development, your baby will receive from 31 to 33 doses of complex biochemical vaccine ingredients. We identify some of these substances, such as adjuvants, preservatives and other 'clinically significant' ingredients in the following popular vaccine brands.

Example Vaccines received in the first year

3 DTaP-HB-IPV-Hib—Infanrix hexa®

2 MEN-C-C—Menjugate[®]
2 Rotavirus—RotarixTM
2 Influenza—Flulaval[®]Tetra
1 MMR—MMRII[®]
1 Varicella—Varivax[®]III

Infanrix hexa:

Adjuvant: Aluminum 820mcg X 3 doses = **2460mcg**

Preservative: 2-Phenoxyethanol (PE)

Potential Allergens: Polymyxin B, Yeast protein. Neomycin,

Tetanus toxoid carrier protein

Other: Bovine serum albumin Formaldehyde Monkey kidney cell DNA Polysorbate 80

Menjugate:

Adjuvant: Aluminum 1000mcg X 1 dose = **1000mcg**

Potential Allergens: Latex, Diphtheria toxoid carrier protein Other: Disodium phosphate heptahydrate, Mannitol, Sodium

chloride, Sodium dihydrogen phosphate monohydrate

Prevnar-13:

Adjuvant: Aluminum 1250mcg X 3 doses = **3750mcg**

Potential Allergens: Diphtheria toxoid carrier protein Other: Polysorbate 20 and 80, Formaldehyde, Lactose

Rotarix: Other: DNA fragments from porcine circovirus 1 Contaminants from cell growth medium

Flulaval Tetra:

Preservative: Thimerosal <25mcg mercury X 2doses = **<50mcg**

Potential Allergens: Egg protein, Thimerosal

Other: Formaldehyde

MMRII:

Potential Alergens: Neomycin, Phenol red, Porcine gelatin

Residual components of chick embryo cell cultures

Other: Fetal bovine serum, Human diploid cells (fetal cells)

Contaminants from cell growth medium,

Recombinant human albumin

Monosodium L-glutamate monohydrate

Varivax III:

Potential allergens: Porcine gelatin, fetal bovine serum, neomycin Other: Human diploid cells (fetal cells), monosodium glutamate

Note: Due to space considerations, chemical salts, sugars and many other ingredients are not listed in the Other Ingedients category above. For complete ingredient listings and information on vaccines used in BC, see the BC Centre for Disease Control manual: Communicable Disease Control/Immunization Program/Section VII—Biological Products.

Using the above vaccines in the first year of life, vaccinated babies will receive a total of 7210 micrograms (mcg) of aluminum and nearly 50 micrograms of mercury, both of which are known neurotoxins affecting brain development and function. Babies will also be exposed to various animal and human DNA, cell fragments and contaminants. The polysorbate 20 & 80 emulsifier can cross the blood brain barrier, ferrying any other ingredients into the brain.

Ask your doctor what vaccines they use and then read the eyeopening product monographs. <u>Product monographs</u> are available online at www.vaccines411.ca under the health professionals tab.

BC Schedule VCC May 2017