








compares Yukon Vaccine Schedules

1983 TOTAL PreSchool 22 Vaccines In 6 Injections & 4 oral doses	AGE Prenatal	2017 TOTAL PreSchool 50 Vaccines In 20 Injections & 2 oral doses
∅ Number Vaccines		1 Influenza Vaccine in routine schedule for pregnant women
DPT: 3 vaccines in 1 shot OPV: oral polio, live virus vaccine 4	2 Months 	9 DTaP-HB-IPV-Hib 6 vaccines in 1 shot Meningococcal C-C Pneumococcal C-13 Rotavirus (oral) live virus vaccine
DPT: 3 vaccines in 1 shot OPV: oral polio live virus vaccine 4	4 Months 	8 DTaP-HB-IPV-Hib 6 vaccines in 1 shot Pneumococcal C-13 Rotavirus (oral) live virus vaccine
DPT: 3 vaccines in 1 shot 3 ∅	6 Months 	8 DTaP-HB-IPV-Hib 6 vaccines in 1 shot Influenza 2 doses 4 weeks apart
∅ 3 MMR: 3 live virus vaccines in 1 shot at 15 months	12 Months 	6 MMR 3 live virus vaccines in 1 shot Meningococcal C-C Pneumococcal C-13 Varicella chickenpox live virus vaccine
DPT: 3 vaccines in 1 shot OPV: oral polio live virus vaccine 4	18 Months 	6 DTaP-IPV-Hib 5 vaccines in 1 shot Influenza (1st Annual dose)
∅ 4 DPT: 3 vaccines in 1 shot OPV: oral polio live virus vaccine Total 22	3 Yr 4-6 Yr 	1 Influenza (Annual dose in Fall) 11 Tdap-IPV 4 vaccines in 1 shot MMRV 4 live virus vaccines in 1 shot Influenza (3 Annual doses, 1 each Fall) Total 50



By the time your child graduates from high school, they will have received

- an additional 11 annual Influenza vaccines,
- 2 doses HPV, girls only in Grade 6
- Tdap (3-in-1 shot) and Men C-ACYW-135 in Grade 9

TOTAL Grade School & High School in 2017

Girls: 17 Vaccines in 15 Injections
Boys: 15 Vaccines in 13 Injections

2017 GRAND TOTAL: Prenatal to Age 18
65 or 67 Vaccines

Did you know vaccination is VOLUNTARY in Canada?

As Health Canada explained years ago: “Unlike some countries, immunization is not mandatory in Canada; it cannot be made mandatory because of the Canadian Constitution...legislation and regulations must not be interpreted to imply compulsory immunization.”

—Immunization in Canada, Volume: 23S4, May 1997, [Canadian National Report on Immunization](#)

LEGEND

Vaccines used in the 1983 Yukon schedule

Note: All 3 of these vaccines have since been withdrawn from the market due to safety concerns.

DPT: diphtheria, whole cell pertussis, tetanus vaccine
3 vaccine combination

OPV: oral, live virus polio vaccine

MMR: measles, mumps (Urabe strain), rubella vaccine
3 live virus vaccine combination

Vaccines used in the 2017 Yukon schedule

DTaP-HB-IPV-Hib 6 vaccine combination with diphtheria(D), tetanus(T), acellular pertussis(aP), HepatitisB (HB), inactivated polio (IPV) & Hib (Haemophilus Influenzae B)
• **INFANRIX** hexa[®]

DTaP-IPV-Hib 5 vaccine combination with diphtheria(D), tetanus(T), acellular pertussis(aP), inactivated polio (IPV) & Hib (Haemophilus Influenzae B) vaccines
• **Pediacel**[™] or **Infanrix-IPV/Hib**[®]

HPV Human papillomavirus vaccine, Quadrivalent
• **GARDASIL** 4[™] (girls only)

Influenza Annual Flu shot, quadrivalent vaccine
• **FLULAVAL**[®]Tetra for babies 6 –23 months
• **FLUMIST**[®] Quadrivalent or **FLULAVAL**[®] Tetra 2 –17 yrs

Men C-C: Meningococcal serotype C vaccine,
• **MENJUGATE**[®] or **NEISVAC-C**[®]

Men C-ACYW-135: 4 Meningococcal serotypes,
• **MENVEO**[®] or **MENACTRA**[®]

***MMR:** measles, mumps, rubella,
3 combination **live virus** vaccine
• **MMRII**[®] or **Priorex**[®] for children < 4 years old

***MMRV:** measles, mumps, rubella, & varicella (chickenpox)
4 combination **live virus** vaccine
• **Priorex Tetra**[®] or **ProQuad**[™] for children ≥ 4 yrs old

Pneu C-13: pneumococcal vaccine, 13 serotypes,
• **PREVNAR** 13[®]

***Rotavirus:** **live virus** oral vaccine,
• **Rotarix**[™] (2 doses) or **RotaTeq**[®] (3 doses)

Tdap-IPV: tetanus, diphtheria, acellular pertussis, polio
4 vaccine combination, age 4-6
• **ADACEL-Polio**[®] or **BOOSTRIX-Polio**[®]

Tdap: tetanus, diphtheria, acellular pertussis, polio
4 vaccine combination, Grade 9 booster
• **ADACEL**[®] or **BOOSTRIX**[®]

***Varicella:** chickenpox **live virus** vaccine
• **VARILRIX**[®] or **VARIVAX**[®] III

***Live virus vaccines** can shed for varying amounts of time (weeks to months) in the body fluids of vaccinated people and can be transmitted to others. *“Little is known about the potential of live attenuated and genetically engineered vaccine viruses to mutate and recombine with other viruses and create new viruses that will cause disease or affect the integrity of the human genome, human microbiome and healthy functioning of the immune and neurological systems.”* **NVIC** <http://www.nvic.org/cmstemplatess/nvic/pdf/live-virus-vaccines-and-vaccine-shedding.pdf>

What’s In those Vaccines?

In the first year of life, during critical phases of brain and immune system development, your baby will receive 32 doses of complex biochemical vaccine ingredients. We identify some of these substances, such as adjuvants, preservatives and other ‘clinically significant’ ingredients in the following popular vaccine brands.

Example Vaccines received in the first year

3 DTaP-HB-IPV-Hib—Infanrix hexa[®]

2 MEN-C-C—Menjugate[®]

2 Rotavirus—Rotarix[™]

1 MMR—MMRII[®]

3 Pneu C-13—Prevnar 13[®]

2 Influenza—Flulaval[®]Tetra

1 Varicella—Varivax[®]III

Infanrix hexa:

Adjuvant: Aluminum 820mcg X 3 doses = **2460mcg**
Preservative: 2-Phenoxyethanol (PE)
Potential Allergens: Polymyxin B, Yeast protein, Neomycin, Tetanus toxoid carrier protein
Other: Bovine serum albumin Formaldehyde
Monkey kidney cell DNA Polysorbate 80

Menjugate:

Adjuvant: Aluminum 1000mcg X 2 doses = **2000mcg**
Potential Allergens: Latex, Diphtheria toxoid carrier protein
Other: Disodium phosphate heptahydrate, Mannitol, Sodium chloride, Sodium dihydrogen phosphate monohydrate

Prevnar-13:

Adjuvant: Aluminum 1250mcg X 3 doses = **3750mcg**
Potential Allergens: Diphtheria toxoid carrier protein
Other: Polysorbate 20 and 80, Formaldehyde, Lactose

Rotarix: Other: DNA fragments from porcine circovirus 1
Contaminants from cell growth medium

Flulaval Tetra:

Preservative: Thimerosal <25mcg **mercury** X 2doses = **<50mcg**
Potential Allergens: Egg protein, Thimerosal
Other: Formaldehyde

MMRII:

Potential Allergens: Neomycin, Phenol red, Porcine gelatin
Residual components of chick embryo cell cultures
Other: Fetal bovine serum, Human diploid cells (fetal cells)
Contaminants from cell growth medium,
Recombinant human albumin
Monosodium L-glutamate monohydrate

Varivax III:

Potential allergens: Porcine gelatin, fetal bovine serum, neomycin
Other: Human diploid cells (fetal cells), monosodium glutamate

Note: Due to space considerations, chemical salts, sugars and many other ingredients are not listed in the Other Ingredients category above. For complete ingredient listings and information on vaccines used in YK, see the YK Health & Social Services, *Immunization Program Manual: Section 8—Biological Products.*

Using the above vaccines, in the first year of life vaccinated babies will receive a total of **8210 micrograms (mcg)** of aluminum and nearly **50 micrograms of mercury**, both of which are known neurotoxins affecting brain development and function. Babies will also be exposed to various animal and human DNA, cell fragments and contaminants. The polysorbate 20 & 80 emulsifier can cross the blood brain barrier, ferrying any other ingredients into the brain.

Ask your clinic what vaccines they use and then read the eye-opening product monographs. [Product monographs](#) are available online at www.vaccines411.ca under the health professionals tab.

The Yukon’s Immunization Program is offered through community health centres and in-school programs. Prior to school based clinics, the Community Health Centre will send out consent forms.