



Compare Canada's Vaccine Schedules

Birth to 18 Years Provincial/Territorial Averages

1950s

14 Doses
of 5 Vaccines

- 4 DOSES of Diphtheria
- 4 DOSES of Tetanus
- 4 DOSES of Pertussis
- 1 DOSE of Polio (1957)
- 1 DOSE of Smallpox

1983

23 Doses
of 7 Vaccines

- 5 DOSES of Diphtheria
- 5 DOSES of Tetanus
- 5 DOSES of Pertussis
- 4 DOSES of Polio (Oral)
- 2 DOSES of Rubella
- 1 DOSE of Measles
- 1 DOSE of Mumps

2020

68 or 70 Doses
of 16 or 17 Vaccines

- 1 DOSE of Meningococcal C-ACYW
- 2 DOSES of Meningococcal C-C
- 2 DOSES of HPV
- 2 DOSES of Varicella (Chickenpox)
- 2 DOSES of Rubella
- 2 DOSES of Mumps
- 2 DOSES of Measles
- 3 DOSES of Rotavirus
- 3 DOSES of Hepatitis A
(Aboriginal Only)
- 3 DOSES of Hepatitis B
- 3 DOSES of Pneumococcal C-13
- 4 DOSES of Hib
- 5 DOSES of Polio (IPV)
- 6 DOSES of Pertussis
- 6 DOSES of Tetanus
- 6 DOSES of Diphtheria
- 18 DOSES of Influenza

*Should children
get a shot they don't need?*

*This may be the most important decision
you will ever make for your child.*

Make an informed decision.

Sub-standard Vaccine Safety Testing

No childhood vaccine product licensed for use in Canada or the USA has been tested using the long-term, double blind, placebo-controlled studies required for all other pharmaceutical products. The medical industry uses the monitoring of adverse events following vaccination as the primary method to evaluate safety. This means that our children are injected with products where safety is determined by the amount of injury or death reported after vaccination.

